


Thyroid Cancer and Future Fertility

Thyroid Cancer and Future Fertility



Thyroid Cancer and future fertility

Thyroid cancer is diagnosed in 45,000 individuals each year in the US. Its treatment may affect future fertility in men and women. It is more common in women with female to male ratio of 3 to 1. It is the most rapidly rising cancer in women living in the US. Thyroid cancers are commonly diagnosed in young women in their reproductive years. Treatment of thyroid cancer generally yields excellent results, with the majority of women surviving 10 years or more after diagnosis. Some women develop thyroid cancer due to iodine deficiency in diet or prior neck radiation. Some types of thyroid cancers are related to inheriting an abnormal gene.

Several types of thyroid cancer are recognized 1. Papillary cancer 2. Follicular cancer 3. Medullary cancer 4. Anaplastic cancer 5. Thyroid lymphoma. Papillary and follicular cancers are less invasive tumors and are encountered in the majority

of women diagnosed with thyroid cancer. They also respond to estrogen as they carry estrogen receptors. Estrogen may promote growth of thyroid cancer cells. Thyroid cancers are usually suspected on neck examination followed by ultrasound or Iodine scan then biopsy. In general, treatment of thyroid cancer require total thyroidectomy-surgical removal of the thyroid gland followed by radioactive iodine to ablate any thyroid remnants. This is followed by long term thyroid hormone replacement. Long term follow up is required after treatment.

Effect of thyroid cancer treatment on the ovary

Thyroidectomy followed by thyroid hormone replacement is not known to affect future fertility in men and women. Radioactive iodine can affect the number and quality of eggs remaining in the ovary. The effect is dependent on the dose of radioactive iodine and the age at treatment. Twenty to 30% of women experience transient amenorrhea or irregular menses starting about 3 months after treatment. Normal menses resume about 6 months later. Permanent ovarian failure is rare but may occur in women at age 40 or older at the time of treatment. Increased incidence of miscarriage is reported in the first year after treatment. With the exception of miscarriages, there is no evidence that exposure to radioiodine affects the outcome of subsequent pregnancies and health of borne children.

Effects of radioactive iodine treatment on the testes

Effect of radioactive iodine treatment may be more severe in men. and is related to the total dose of radioactive iodine received. Transient reduction in testosterone and sperm count may occur but sometimes permanent reduction in sperm count and testosterone levels. Men who received large total dose

sometimes sustain permanent damage to the testes with absence of ejaculated sperm-azospermia. There is no evidence of effects of radioactive iodine on their newborn children, although its advised that men avoid fathering children for 6 months after treatment.

Options for fertility preservation

Men interested in future fertility should consider sperm freezing prior to radioiodine treatment. Women should also consider fertility preservation if they will be treated with radioactive iodine and are older than 35 years. Radioiodine treatment will reduce their ovarian reserve. In addition they will be required to avoid pregnancy for a year or so. Options available for preservation of fertility in women include ovarian stimulation and egg retrieval followed by egg or embryo freezing. Ovarian stimulation can be modified to avoid estrogen exposure during stimulation. Moreover, in familial thyroid cancers, embryos can be genetically tested to avoid transmission of the abnormal gene to children. Men and women diagnosed with thyroid cancer can benefit from consultation with a fertility preservation specialist prior to treatment to discuss effects on gonads and methods to preserve future fertility. Read more at <http://nycivf.org>